arXiv:astro-ph/0505588v4 14 Oct 2005

Relic Gravitational Waves and the Evolution of the Universe

Wen Zhao
Astrophysics Center
University of Science and Technology of China
Hefei, Anhui, China

Abstract

In the inflation models, the relic gravitational waves (RGW) generated in the inflation stage, and evolved in the Universe until now. In the different cosmological evolution models, one can get different gravitational waves power spectrum. In this paper, we give a simple formula to estimate this spectrum in a general cosmological model. From this formula, one can easily find the relation between this power spectrum and the cosmological evolution models. The spectrum includes all the information about the evolution of the scale factor a from inflation to now. So RGW is a more clear fossil, which records the cosmological evolution information, and be a useful complementarity for another fossil−Cosmic Microwave Background Radiation (CMB).

PACS numbers: 98.80.-k 98.80.Es 04.30.-w 04.62.+v
e-mail: wzhao7@mail.ustc.edu.cn

In the inflation expansion cosmic models, the stochastic background of relic gravitational waves can generate from the quantum fluctuate, which is important in cosmology and has been extensively studied in the past[1]. The power spectrum of it, which is observed today, depends not only on the details of the early stage of inflationary expansion, but also on the expansion behavior of the subsequent stages[2, 3], especially, some stages as the pre-big bang stage, the reheating stage, the cosmological QCD transition, the e^+e^- annihilation process, the accelerating expansion process, which we did not know very clearly in physics until now. Their information has also been recorded in the RGW power spectrum. Most of these stages occurred before the CMB photon decoupled, which makes it difficult to study them from CMB. So research on RGW power spectrum is nearly the only very way to study this unknown process of the Universe. In this paper, I will study this power spectrum in a general cosmological model, and find that there is a very simple formula to estimate it. From this formula. one can easily find the relation between the spectrum and the evolution of the Universe. Especially, the spectrum index directly relates to the state equation of matter which dominated the evolution of the Universe at some stage. This gives a simple way to estimate the cosmic evolution process from the RGW power spectrum.

In the spatially flat Robertson-Walker spacetime, the metric which includes the gravitational waves is

$$ds^2 = a^2(\tau)[d\tau^2 - (\delta_{ij} + h_{ij})dx^i dx^j], \tag{1}$$

where τ is the conformal time, the perturbations of spacetime h_{ij} is a 3×3 symmetric matrix. The gravitational wave field is the tensorial portion of h_{ij}, which is transverse-traceless $\partial_i h^{ij} = 0$,$\delta^{ij} h_{ij} = 0$. Since the relic gravitational waves are very weak, $|h_{ij}| \ll 1$, so one needs just study the linearized field equation:

$$\partial_\mu(\sqrt{-g}\partial^\mu h_{ij}(\mathbf{x}, \tau)) = 0. \tag{2}$$

In quantum theory of gravitational waves, the field h_{ij} is a field operator, which is written as a sum of the plane wave Fourier modes

$$h_{ij}(\mathbf{x}, \tau) = \frac{\sqrt{16\pi} l_{Pl}}{(2\pi)^{\frac{3}{2}}} \sum_{\lambda=1}^{2} \int_{-\infty}^{\infty} d^3\mathbf{k} \epsilon_{ij}^{(\lambda)}(\mathbf{k}) \frac{1}{\sqrt{2k}} [a_k^{(\lambda)} h_k^{(\lambda)}(\tau) e^{i\mathbf{k}\cdot\mathbf{x}} + a_k^{\dagger(\lambda)} h_k^{(\lambda)*}(\tau) e^{-i\mathbf{k}\cdot\mathbf{x}}], \tag{3}$$

where $l_{Pl} = \sqrt{G}$ is the Planck's length, the two polarizations $\epsilon_{ij}^{(\lambda)}(\mathbf{k})$, $\lambda = 1, 2$, are symmetric and transverse-traceless $k^i \epsilon_{ij}^{(\lambda)}(\mathbf{k}) = 0$, $\delta^{ij} \epsilon_{ij}^{(\lambda)}(\mathbf{k}) = 0$, and satisfy the conditions $\epsilon^{(\lambda)ij}(\mathbf{k}) \epsilon_{ij}^{(\lambda')}(\mathbf{k}) = 2\delta_{\lambda\lambda'}$, and $\epsilon_{ij}^{(\lambda)}(-\mathbf{k}) = \epsilon_{ij}^{(\lambda)}(\mathbf{k})$, the creation and annihilation operators satisfy $[a_{\mathbf{k}}^{(\lambda)}, a_{\mathbf{k}'}^{\dagger(\lambda')}] = \delta_{\lambda\lambda'} \delta^3(\mathbf{k} - \mathbf{k}')$.

For a fixed wave number $\mathbf{k}$ and a fixed polarization state λ, the wave equation reduces to the second-order ordinary differential equation [3, 4]

$$h_k^{(\lambda)''}(\tau) + 2\frac{a'}{a} h_k^{(\lambda)'}(\tau) + k^2 h_k^{(\lambda)} = 0, \tag{4}$$

where the prime denotes $d/d\tau$, and $k = |\mathbf{k}|$. Since the equation of $h_k^{(\lambda)}(\tau)$ for each polarization is the same, we denote $h_k^{(\lambda)}(\tau)$ by $h_k(\tau)$ in below.

The power spectrum $h(k, \tau)$ of relic gravitational waves is always defined as

$$h(k, \tau) = \frac{4 l_{Pl}}{\sqrt{\pi}} k |h_k(\tau)|. \tag{5}$$

The spectral density parameter $\Omega_g(\nu)$ of gravitational waves is defined through the relation $\rho_g/\rho_c = \int \Omega_g(\nu) \frac{d\nu}{\nu}$, where ρ_g is the energy density of the gravitational waves and ρ_c is the critical energy density. One reads

$$\Omega_g(\nu) = \frac{\pi^2}{3} h^2(k, \tau_H) (\frac{\nu}{\nu_H})^2, \tag{6}$$

where τ_H is the conformal time of now and ν_H is the Hubble frequency of now. Which can not exceed 10^{-6} as required by the rate of the primordial nucleogenesis[5].

In the inflation models, RGW generated from the initial quantum fluctuate during the inflationary stage. In general, in the inflation stage, the scale factor has a form

$$a(\tau) = l_0 |\tau|^{1+\beta(\tau)}, \quad -\infty < \tau < 0, \tag{7}$$

Normally, β is a constant. If $\beta = -2$, the inflation is a simple de Sitter expansion. But recently from observed results of the primordial scalar power spectrum from WMAP[6], one finds the inflation is a very complicated process, here we assume β is a slowly evolving function with time τ. For a given wave number k, it crossed over the horizon at a time τ_i, i.e. when the wave length $\lambda_i = 2\pi a(\tau_i)/k$ is equal to $1/H(\tau_i)$, the Hubble radius at time τ_i. Eq. (7) yields $1/H(\tau_i) = l_0|\tau_i|^{2+\beta}/|1+\beta|$, and for the exact de Sitter expansion with $\beta = -2$ one has $H(\tau_i) = l_0$. Note that a different k corresponds to a different time τ_i. Now choose the initial condition of the mode function $h_k(\tau)$ as[3]

$$|h_k(\tau_i)| = \frac{1}{a(\tau_i)}. \tag{8}$$

Then the initial amplitude of the power spectrum is

$$h(k, \tau_i) = 8\sqrt{\pi} l_{pl} H(\tau_i), \tag{9}$$

From $\lambda_i = 1/H(\tau_i)$ it follows that $\frac{a'(\tau_i)}{a(\tau_i)} = \frac{k}{2\pi}$. So the initial amplitude of the power spectrum is

$$h(k, \tau_i) = Ak^{2+\beta(\tau_i)} = Ak^{2+\beta(-1/k)}, \tag{10}$$

where the constant

$$A = 8\sqrt{\pi} l_{pl} |1 + \beta(\tau_i)|/l_0. \tag{11}$$

For the case of $\beta = -2$, the initial spectrum is independent of k, this is the so called scale independent initial power spectrum. Its amplitude directly relate the inflation Hubble constant.

The evolution of RGW in the following stage satisfies the two order differential equation (4). Here we only suppose that the Universe is always a decelerating expand. (from the research of SNIa, WMAP and LSS, one finds that now the Universe is accelerating expansion, we have researched its effect on the spectrum and found it was little[3].) If the matter state equation satisfies $p = \omega\rho$, where p and ρ are the pressure and energy density of the matter respectively. (ω is a constant, for radiation, $\omega = 1/3$; and for matter $\omega = 0$), one can easily find that the scale factor has a form:

$$a = b[\tau - \tau_1]^{\alpha}, \tag{12}$$

where $\alpha = \frac{2}{3\omega+1}$, b and τ_1 are constant. For the component of the Universe vary with time, we here think the ω is varying slowly with time, this is a better way to describe the evolution of the Universe than the method which separated the Universe into several different evolution stages. So the scale factor can be described in the following form:

$$a = b(\tau)[\tau - \tau_1(\tau)]^{\alpha(\tau)}, \tag{13}$$

where $b(\tau)$, $\tau_1(\tau)$ and $\alpha(\tau)$ are all slowly evolution functions with time. This way can be used to study all kind of cosmic evolution models. Especially, it is useful for studying the effect of the cosmological phase transition on the RGW power spectrum.

In this model, when the wavelength of RGW is much larger than the horizon, the amplitude of the spectrum keeps constant. Once the gravitational waves enters the horizon, the spectrum will have a damping along with the expansion the Universe. We assume that at a certain short stage, the scale factor has a form of

$$a = b'[\tau - \tau_1']^{\alpha'}, \tag{14}$$

For $b'(\tau)$, $\tau_1'(\tau)$ and $\alpha'(\tau)$ vary slowly with time, they can be assumed as constant in this stage. We study the gravitational waves which enter the horizon in this stage. A wave with wavenumber k, its amplitude satisfies[3]

$$h(\tau_H, k) = h(\tau_i, k) a_*(\tau)/a(\tau_H), \tag{15}$$

where $a_*(\tau)$ is the scale factor when the wave k just enters horizon, and $a(\tau_H)$ is the now scale factor. From the scale factor formula and the relation $k(\tau - \tau_1') \simeq 1$ when wave enters horizon one find

$$a_*(\tau)/a(\tau_H) = \frac{b(1/k)}{b_H} \frac{k_H^{\alpha(\tau_H)}}{k^{\alpha}}, \tag{16}$$

where $b_H \equiv b(\tau_H)$. So we can get the spectrum

$$h(k, \tau_H) = (A/b_H) b(1/k) k_H^{\alpha(\tau_H)} k^{\beta(-1/k)+2-\alpha(1/k)} = 8\sqrt{\pi} l_{pl} H(\tau_i)(b(1/k)/b_H) k_H^{\alpha(\tau_H)} k^{-\alpha(1/k)}, \tag{17}$$

For the slowly varying with time of $b(\tau)$, $\tau_1(\tau)$ and $\alpha(\tau)$, we can use this formula to estimate the RGW spectrum for all k. From this, one can also get the spectrum index

$$n(k) \equiv d \ln h(k, \tau_H)/dk \simeq \beta(-1/k) + 2 - \alpha(1/k), \tag{18}$$

So the matter state equation when the horizon of the Universe equate to $1/k$ is

$$\omega(\tau) = \omega(1/k) = \frac{2}{3[\beta(-1/k) + 2 - n(k)]} - 1, \tag{19}$$

In general, $\beta + 2$ is nearly zero for inflation, so this formula directly relates the RGW spectrum index and matter state equation in the Universe. For the waves with wavelength long enough to have not reenter the horizon until now, they have the power spectrum of their initial condition

$$h(k, \tau_H) = h(k, \tau_i) = 8\sqrt{\pi} l_{pl} H(\tau_i) = A k^{2+\beta(-1/k)}, \tag{20}$$

and

$$n(k) \simeq \beta(-1/k) + 2. \tag{21}$$

These are the approximate formulae, which can evaluate the RGW power spectrum (If one want to know the exact spectrum, it must depend on the numerical calculation or the complicated exact calculation), but it is enough for knowing the physical information which included in the RGW power spectrum. From formula (20), one can find these spectrum keeps its initial value, which reflects the physics of the inflation. Its amplitude is exactly the Hubble parameter of the inflation, its change with k is also the change of Hubble parameter with time. So from which we can know not only the energy scale of the inflation which closely relates the inflation physics origin, but also the inflation evolution information, which is also very important for the very damping of the CMB anisotropies power spectrum in the very large scale, and the large running of index of the primordial scalar perturbation observed by WMAP[6, 7], some people think that it reflects the complicated evolution of early inflation.

For the waves with larger frequency, the spectrum formula (17) also includes the evolution information of the Universe after inflation. Its amplitude is dependent on the all evolution process of Universe after cosmic horizon scale became larger than the wavelength. Especially the spectrum index n_s, which directly relates to the matter state equation when the cosmic horizon scale equate to the wavelength, and the variety of the spectrum index with wavelength is also the variety of the matter state equation in the Universe with time. Actually, we can only detect the RGW power spectrum in several narrow frequency span, for example the laser interference explorers[8], which is sensitive for the wave with frequency of $10^{-3} \sim 10^{2}$Hz, the method using the millisecond pulsars[9], which is sensitive at $10^{-9} \sim 10^{-7}$Hz, the method by researching the anisotropy and polarization of CMB[10], which is sensitive at $10^{-18} \sim 10^{-16}$Hz. So even if we have detected RGW at some frequency, we also can't know clearly the inflation energy scale unless we know all information of Universe after the inflation stage, but we can research the matter state equation in a special stage of the Universe by the index of the power spectrum, which directly relates the matter state equation ω at the time $\tau = 1/k$ by formula (19). So I think the simple RGW power spectrum in all frequency can reflect the all evolution information of the Universe since inflation, and it can not be contaminated by other information as matter perturbation, the interaction between the matters. So it is a very clear fossil for study the cosmic evolution, and be a useful complementarity for another fossil−CMB, especially for studying of the Universe stage before the CMB photon decoupled.

References

[1] S.Sasaki, Prog.Theor.Phys. **76**,1036 (1986); A.A.Starobinsky, JEPT Lett. **30**, 682 (1979); A. Rubakov, M.V.Sazhin, and A.V.Veryaskin, Phys.Lett.**115B**, 189 (1982); R.Fabbri and M.D. Pollock, Phys.Lett. **125B**, 445 (1983); L.F.Abbott and M. B. Wise, Nucl.Phys. **B224**, 541 (1984); L.F.Abbott and D.D.Harari, Nucl.Phys. **B264**, 487 (1986); V.F.Mukhanov, H.A.Feldman, and R.H.Brandenberger Phys.Report **215**,203 (1992); D. H. Lyth, A. Riotto, Phys.Rept. **314** 1 (1999);

[2] B.Allen, Phys.Rev.D **37**, 2078 (1988); V.Sahni Phys.Rev.D **42**, 453 (1990); L.Grishchuk, Class.Quant.Grav. **14** 1445 (1997); Lecture Notes Physics **562**, 164 (2001), in "*Gyros, Clocks, Interferometers...:Testing Relativistic Gravity in Space*", Lammerzahl et al (Eds); A. Riszuelo and J-P Uzan, Phys.Rev.D 62, 083506 (2000); H. Tashiro, K. Chiba, and M. Sasaki, Class.Quant.Grav. **21** 1761 (2004); A.B. Henriques, Class.Quant.Grav. **21** 3057 (2004);

[3] Y.Zhang, Y.F.Yuan, W.Zhao, Y.T.Chen, Class.Quant.Grav. **22**, 1383 (2005); Y.Zhang, W. Zhao, Y.F. Yuan and T.Y. Xia Chin.Phys.Lett. **20**, 1871 (2005);

[4] L.Grishchuk, Sov.Phys.JETP, Vol. **40**, No.3, 409 (1974).

[5] E.Kolb and R.Turner, *The Early Universe* (Addison-Wesley, Reading, MA, 1990); B.Allen, gr-qc/9604033; M.Maggiore, Phys.Rep. **331**, 283 (2000);

[6] C.L.Bennett, et al. Astrophys.J.Suppl. **148**, 1 (2003); D.N.Spergel, et al. Astrophys.J.Suppl. **148**, 175 (2003);

[7] G. Hinshaw, et al. Astrophys.J.Suppl. **148**, 135 (2003); H.V.Peiris, et al. Astrophys.J.Suppl. **148**, 213 (2003);

[8] http: // www.ligo.caltech.edu; http://www.geo600.uni.hannover.de; http://http://lisa.jpl.nasa.gov/; LIGO Scientific Collaboration: B. Abbott, et al, astro-ph/0507254;

[9] S. Detweiler, Astrophys. J. **234**, 1100 (1979); A.N. Lommen, astro-ph/0208572;

[10] M. Kamionkowski and A. Kosowsky, Ann. Rev. Nucl. Part. Sci. **49**, 77 (1999); W. Hu and S. Dodelson, Ann. Rev. Astron. Astrophys. **40**, 171 (2002); M. White and J. D. Cohn, astro-ph/0203120; S. Dodelson, *Modern Cosmology* (Academic Press, 2003); A. Liddle and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure* (Cambridge University Press, 2000); Wen Zhao and Yang Zhang, astro-ph/0509753, astro-ph/0510355;